



Mokaya, J., Maponga, T. G., McNaughton, A. L., Van Schalkwyk, M., Hugo, S., Singer, J. B., Sreenu, V. B., Bonsall, D., de Cesare, M., Andersson, M., Gabriel, S., Taljaard, J., Barnes, E., Preiser, W., Van Rensburg, C., & Matthews, P. C. (2020). Evidence of tenofovir resistance in chronic hepatitis B virus (HBV) infection: An observational case series of South African adults. *Journal of Clinical Virology*, 129, 104548. <https://doi.org/10.1016/j.jcv.2020.104548>

Publisher's PDF, also known as Version of record

License (if available):  
CC BY

Link to published version (if available):  
[10.1016/j.jcv.2020.104548](https://doi.org/10.1016/j.jcv.2020.104548)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the final published version of the article (version of record). It first appeared online via Elsevier at <https://doi.org/10.1016/j.jcv.2020.104548> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>



## Short communication

## Evidence of tenofovir resistance in chronic hepatitis B virus (HBV) infection: An observational case series of South African adults

Jolynne Mokaya<sup>a,1</sup>, Tongai G Maponga<sup>b,1</sup>, Anna L McNaughton<sup>a</sup>, Marije Van Schalkwyk<sup>c</sup>, Susan Hugo<sup>c</sup>, Joshua B Singer<sup>d</sup>, Vattipally B Sreenu<sup>d</sup>, David Bonsall<sup>a,e,f</sup>, Mariateresa de Cesare<sup>g</sup>, Monique Andersson<sup>b,e</sup>, Shiraaz Gabriel<sup>h</sup>, Jantje Taljaard<sup>c</sup>, Eleanor Barnes<sup>a,i,j</sup>, Wolfgang Preiser<sup>b</sup>, Christo Van Rensburg<sup>h</sup>, Philippa C Matthews<sup>a,e,j,\*</sup>

<sup>a</sup> Nuffield Department of Medicine, University of Oxford, Medawar Building, South Parks Road, Oxford OX1 3SY, UK

<sup>b</sup> Division of Medical Virology, Stellenbosch University / National Health Laboratory Service Tygerberg, Cape Town, South Africa

<sup>c</sup> Division of Infectious Diseases, Department of Medicine, Stellenbosch University / Tygerberg Academic Hospital, Cape Town, South Africa

<sup>d</sup> MRC-University of Glasgow Centre for Virus Research, Bearsden Road, Glasgow, G61 1QH, UK

<sup>e</sup> Department of Infectious Diseases and Microbiology, Oxford University Hospitals NHS Foundation Trust, John Radcliffe Hospital, Headley Way, Oxford OX3 9DU, UK

<sup>f</sup> Big Data Institute, Old Road, Oxford OX3 7FZ, UK

<sup>g</sup> Wellcome Centre for Human Genetics, Roosevelt Drive, Oxford OX3 7BN, UK

<sup>h</sup> Division of Gastroenterology, Department of Medicine, Stellenbosch University / Tygerberg Academic Hospital, Cape Town, South Africa

<sup>i</sup> Department of Hepatology, Oxford University Hospitals NHS Foundation Trust, John Radcliffe Hospital, Headley Way, Oxford OX3 9DU, UK

<sup>j</sup> National Institutes of Health Research Health Informatics Collaborative, NIHR Oxford Biomedical Research Centre, John Radcliffe Hospital, Headley Way, Oxford OX3 9DU, UK

## ARTICLE INFO

## Keywords:

HBV  
HIV  
Tenofovir  
TDF  
Resistance  
RAMs  
Africa  
Nucleos(t)ide analogue  
Therapy  
Viraemia

## ABSTRACT

**Introduction:** Tenofovir disoproxil fumarate (TDF) is widely recommended for treatment of chronic hepatitis B virus (HBV) infection because it is safe, affordable and has a high genetic barrier to resistance. TDF resistance associated mutations (RAMs) have been reported, but data are limited, particularly for Africa. We set out to identify potential RAMs in individuals with detectable HBV viraemia on TDF treatment.

**Methods:** We recruited adults with chronic HBV infection from Cape Town, South Africa, identifying individuals with a TDF resistance phenotype, defined as persistent HBV viraemia despite > 12 months of TDF treatment. We sequenced HBV DNA using MiSeq Illumina with whole genome target enrichment, and sought potential TDF RAMs, based on a pre-defined list of polymorphisms.

**Results:** Among 66 individuals with chronic HBV (genotypes A and D), three met our clinical definition for TDF resistance, of whom two were coinfecting with HIV. In one participant, the consensus HBV sequence contained nine polymorphisms that have been described in association with TDF resistance. Significant treatment non-adherence in this individual was unlikely, as HIV RNA was suppressed. TDF RAMs were also present in HBV sequences from the other two participants, but other factors including treatment non-adherence may also have had a role in failure of HBV DNA suppression in these cases.

**Discussion:** Our findings add to the evidence that RAMs in HBV reverse transcriptase may underpin a TDF resistant phenotype. This is the first time these RAMs have been reported from Africa in association with clinical evidence of TDF resistance.

## 1. Introduction

Tenofovir disoproxil fumarate (TDF) is a first line nucleoside analogue (NA) agent used to treat chronic hepatitis B virus (HBV) infection [1,2]. Although the genetic barrier to TDF resistance is high [1,2],

recent reports suggest possible HBV resistance to TDF [3,4]. A review of potential TDF resistance associated mutations (RAMs) highlights significant gaps and limitations in the current literature, demonstrating the need for further investigation [5].

TDF is also used as antiretroviral therapy (ART) for HIV [6,7]. Thus,

\* Corresponding author at: Nuffield Department of Medicine, Medawar Building, South Parks Road, Oxford OX1 3SY, UK.

E-mail address: [philippa.mathews@ndm.ox.ac.uk](mailto:philippa.mathews@ndm.ox.ac.uk) (P.C. Matthews).

<sup>1</sup> These authors contributed equally to this article.

in settings where HIV is widespread, population exposure to TDF is high, raising the potential risk of selection of resistance in both HIV and HBV [7]. Monitoring HIV viraemia in coinfecting patients receiving TDF-based regimens provides insights about treatment compliance, as HIV suppression is a surrogate for adherence to therapy.

Improved insight into the prevalence and clinical significance of HBV RAMs is essential; selection and dissemination of TDF resistant HBV would limit the success of antiviral therapy for individual patients, and may undermine efforts to curb transmission in pursuit of elimination targets [8–10]. We identified patients in whom concerns about possible drug resistance were raised on account of persistent HBV viraemia despite TDF therapy. Sequencing HBV confirmed the presence of mutations that have previously been described in association with TDF resistance.

## 2. Methods

### 2.1. Clinical cohort

Adults were enrolled into a cross-sectional cohort of CHB at Tygerberg Hospital, Cape Town, South Africa ('OxSAHep' HBV study), as previously described [11]. All participants provided written informed consent. Ethics approval was provided by the University of Oxford Tropical Research Ethics Committee (ref. OXTREC 01–18) and Stellenbosch University Human Research Ethics Committee (HREC ref. N17/01/013). We identified patients with CHB in whom HBV viraemia was  $\geq 20,000$  IU/mL despite being on TDF treatment for  $> 12$  months (Fig. 1).

South African guidelines for HBV treatment recommend use of lamivudine (3TC), entecavir (ETV) or TDF [12,13]. In practice, TDF is most frequently used, as 3TC use is limited by drug resistance, and ETV is expensive and not widely available.

### 2.2. Deep sequencing and identification of RAMs

We extracted HBV DNA from plasma using a NucliSENS magnetic extraction system (BioMérieux), and used a published protocol to generate fully double stranded HBV genomes [14,15]. We purified DNA and prepared sequencing libraries followed by whole genome target-enrichment with custom-designed HBV probes. We sequenced the samples on an Illumina Mi-Seq using a v3 300bp paired end kit and analysed using Tanoti v1.3 [16] with genotyping and variant calling

performed using the HBV-GLUE resource [17]. We called TDF RAMs using a catalogue of 37 RAMs in HBV reverse transcriptase (RT) [5,7].

## 3. Results

### 3.1. Phenotypic evidence of TDF resistance

Among 66 adults with CHB, we identified three with HBV DNA viral load (VL)  $\geq 20,000$  IU/mL, despite therapy with TDF for  $\geq 12$  months (Figure 1), of whom two were co-infected with HIV. Both co-infected patients were treated with 3TC and Emtricitabine (FTC) in addition to TDF. In patient 209, HIV RNA suppression provides evidence for treatment adherence (Figure 2A Figure 2A), increasing the probability that HBV drug resistance is genuinely present. In contrast, in patient 258, both HIV and HBV remained detectable, suggesting a possible incomplete adherence to therapy, with or without drug resistance in one or both viruses (Figure 2B Figure 2B).


A third case (ID 289) was HIV negative. HBV viral suppression occurred after a cumulative 30 months on treatment (Figure 2C Figure 2C). This individual also had concurrent large flares in liver enzymes (Figure 2 C Figure 2C).

### 3.2. Genomic evidence of TDF resistance


We derived full-length HBV genome sequences for all three patients, although with varying depth of coverage, identifying genotype-A in two cases and -D in the third (Table 1). The consensus sequences all contained RAMs that have been previously described in association with TDF resistance.

In HBV sequences from patient 209, we observed nine potential TDF RAMs. Among these, four polymorphisms (R153W, V173L, L180M, and M204V), have the strongest evidence base [5]. M204I/V and L180M are best recognised in association with 3TC resistance [19], but they may also be part of a combination of polymorphisms that underpins TDF resistance as there was no sustained viral suppression. In patients 258 and 289, six and five putative TDF RAMs were observed, respectively. At a deep sequence level, the RAMs we observed in all three patients were present in  $> 95\%$  of HBV reads.

For each potential RAM identified in our HBV sequences we reviewed the prevalence of the variants in published HBV sequences (Table 2). In all three cases, a number of potential TDF RAMs actually represent the most common residue in this genotype [20]. Of note, for



**Fig. 1.** Selection of participants with chronic HBV infection for viral sequencing to identify tenofovir disoproxil fumarate (TDF) resistance associated mutations (RAMs). Adults attending clinical follow-up in Cape Town, South Africa, were enrolled over a period of one year. Three adults with persistent viraemia despite  $> 12$  months of TDF therapy are indicated in the blue box. Of these one had a phenotypic profile and HBV sequence data to support TDF resistance (orange box); two others had possible TDF resistance RAMs in HBV sequences but other factors including non-adherence to therapy may have contributed to persistent viraemia (yellow box). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article).



**Fig. 2.** Longitudinal viral load and liver function test distribution of three adults with detectable HBV viraemia > 20,000 IU/mL despite ≥12 months prescribed TDF therapy. Panels A-C plot laboratory parameters over time in three patients. The corresponding clinical data for each patient is shown below in panel D.

**Table 1**

Summary of patient characteristics and HBV sequence data generated from three South African adults. All patients had with persistent HBV viraemia ≥20,000 IU/mL despite > 12 months prescription of TDF therapy.

| Patient ID  | 209  | 258                                     | 289  |
|---|--|---|--|
| Treatment   | TDF + 3TC + FTC  | TDF + 3TC + FTC                         | TDF  |
| Age range at recruitment  | 41–50  | 41–50                                   | 61–70  |
| HIV status  | Positive   | Positive                                | Negative                                     |
| Elastography score (kPa)  | 6.9  | 5.1                                     | 9.1  |
| HBV DNA viral load, log <sub>10</sub> IU/mL at time of sequencing     | 8.03   | 6.77                                    | 8.18   |
| HIV RNA viral load, log <sub>10</sub> copies/mL at time of sequencing | 1.74   | 4.97                                    | N/A  |
| Median coverage/site  | 4262   | 1601                                    | 14   |
| HBV Genotype  | A  | D                                       | A  |
| Reverse transcriptase (RT) polymorphisms                              | <u>Y9H</u><br><u>L91I</u>  | <u>Y9H</u><br><u>F122L</u>              | <u>Y9H</u><br><u>L91I</u>                    |
| potentially associated with TDF resistance                            | <u>H126Y</u><br>R153W*<br>V173L <sup>*,†</sup><br>L180M <sup>*,†</sup><br>M204V <sup>*,†</sup><br><u>S223A</u><br><u>C256S</u> | <u>S223A</u><br>D263E<br>V278I<br>A317S | <u>H126Y</u><br><u>S223A</u><br><u>C256S</u> |

Underlined polymorphisms have been reported as RAMs, but occur frequently in the genotype identified in this patient (see Table 2).

\* These RAMs have been identified based on a combination of both *in vivo* and *in vitro* data and are reported in ≥2 independent peer-reviewed sources, giving them the most stringent evidence base of all the polymorphisms listed [6].

† These RAMs cause HBV resistance to 3TC [8]. As patient 209 was on both TDF and 3TC, it is likely that exposure to 3TC resulted in the selection of RAMs V173L, L180M and M204V.

patient 289, four of five TDF RAMs are actually the most common residue at the site for genotype-A.

## 4. Discussion

### 4.1. Summary of findings

We present the first evidence of potential TDF resistance in adults with CHB in Africa. The three patients presented reflect different

**Table 2**

Frequency of putative TDF RAMs in genotype A (n = 290) and D (n = 566) HBV sequences downloaded from Hepatitis B Virus Database (HBVdb), <https://hbvdb.ibcp.fr/HBVdb/>. 840 genotype A and 958 genotype D sequences were downloaded on the 20th of November 2018, and the datasets were filtered to remove highly similar sequences using a 99.5 % similarity threshold, leaving a total of 290 geno-A and 566 geno-D sequences.

| Polymorphism in HBV RT | Frequency in genotype A sequences | Frequency in genotype D sequences |
|------------------------|-----------------------------------|-----------------------------------|
| Y9H                    | 286/290 (98.6 %)                  | 556/566 (98.2 %)                  |
| L91I                   | 290/290 (100 %)                   | 41/566 (7.2 %)                    |
| F122Y/L                | 8/290 (2.8 %)                     | 82/566 (14.5 %)                   |
| H/126Y                 | 225/290 (77.6 %)                  | 4/566 (0.7 %)                     |
| R153W                  | 228/290 (78.6 %)                  | 22/566 (3.9 %)                    |
| V173L                  | 0/290 (0%)                        | 4/566 (0.7 %)                     |
| L180M                  | 7/290 (2.4 %)                     | 13/566 (2.3 %)                    |
| M204I/V                | 7/290 (2.4 %)                     | 25/566 (4.4 %)                    |
| S223A                  | 289/290 (99.7 %)                  | 563/566 (99.5 %)                  |
| C256S                  | 279/290 (96.2 %)                  | 65/566 (11.5 %)                   |
| D263E                  | 1/290 (0.3 %)                     | 111/566 (19.6 %)                  |
| V278I                  | 16/290 (5.5 %)                    | 55/566 (9.7 %)                    |
| A317S                  | 4/290 (1.4 %)                     | 531/566 (93.8 %)                  |

phenotypes, with one case (ID 209, Figure 2A) representing particularly strong evidence for HBV resistance to TDF. The other two cases illustrate the difficulties in discriminating between drug resistance and incomplete adherence, and may also reflect the long timelines sometimes associated with viraemic suppression on TDF [21]. These explanations are not mutually exclusive, and may all have a role in incomplete virologic suppression.

### 4.2. Interpretation

Patient 209 was treated with both TDF and 3TC, making it likely that exposure to 3TC resulted in the development of RAMs V173L, L180M and M204V, possibly setting the scene for the subsequent development of cross-resistance to TDF. Although FTC is not part of the recommended treatment options for CHB, it is widely used for HIV and has some activity against HBV. Combination TDF and FTC has been shown to have better viral suppression compared to TDF monotherapy among treatment naïve individuals with CHB [22], suggesting these mutations may confer resistance to both drugs.

The precise contribution of these polymorphisms to resistance



remains to be determined with certainty [5]. All putative RAMs identified in patient 289 occur frequently in genotype-A sequences, making it uncertain whether drug resistant polymorphisms have been selected *de novo* or are part of a founder sequence. This individual also had large flares in liver enzymes (Figure 2C Figure 2C) that may be evidence of a host immunological response in parallel with drug therapy accounting for the reduction in HBV DNA viraemia. Identifying putative RAMs, and ascertaining their significance to clinical phenotype, is more difficult when these polymorphisms represent the 'wild type' sequence in certain genotypes (as described for hepatitis C virus [24,25]). There is a pressing need for more clinical data combined with *in vitro* data from assays that can confirm a resistant phenotype.

#### 4.3. Caveats and limitations

The data we present here are limited to only three individuals, with different clinical phenotypes (Fig. 2). In patient 289, despite a high viral load we achieved a limited sequencing coverage depth (Table 2). Incomplete treatment adherence can also result in virological breakthrough, and assessment of drug levels in plasma and/or PBMCs could provide more objective information [23], especially where HIV cannot be used as a surrogate measure of treatment compliance.

#### 4.4. Summary

As TDF is widely used in Africa as a component of first line ART, it is possible that the selection of RAMs in HBV is more prevalent than previously reported [7,12]. Our findings add to the evidence that mutations in HBV RT can underpin a TDF resistant phenotype. This is the first time these RAMs have been reported from Africa supported by clinical evidence of TDF resistance. Further investigation is urgently needed to determine the prevalence and clinical significance of TDF RAMs, to investigate the role of combination therapy, and to support the development of new antiviral agents.

#### Data availability

HBV sequence data are available at GenBank [accession numbers MT210032, MT210033, MT210034]

#### Authors contribution statement

JM, TM and PM conceived the study. TM and PM acquired funding and ethics approval. TM, MvS, SH, JT, WP, CvR recruited patients and collected clinical data. JM, AM, DB and MdC generated sequence data. JM, TM, AM, JS vs and PM analysed the data. JM, AM and PM wrote the manuscript. All authors revised and approved the final manuscript.

#### Financial support statement

This work was supported by the Leverhulme Mandela Rhodes Scholarship to JM, Wellcome Trust (grant number 110110Z/15/Z to PCM), the Medical Research Council UK to EB, the Oxford NIHR Biomedical Research Centre to EB. EB is an NIHR Senior Investigator. The views expressed in this article are those of the author and not necessarily those of the NHS, the NIHR, or the Department of Health.

#### Declaration of Competing Interest

We have no conflicts of interest to declare.

#### References

- [1] World Health Organisation, Guidelines for the Prevention, Care and Treatment of

- Persons with Chronic Hepatitis B Infection, Available at: <http://www.who.int/hepatitis/publications/hepatitis-b-guidelines/en/> [Accessed February 22, 2020] (2019).
- [2] P. Marcellin, D.K. Wong, W. Sievert, P. Buggisch, J. Petersen, R. Flisiak, M. Manns, K. Kaita, Z. Krastev, S.S. Lee, et al., Ten-year efficacy and safety of tenofovir disoproxil fumarate treatment for chronic hepatitis B virus infection, *Liver Int.* (2019) 1868–1875.
- [3] W.H. Cho, H.J. Lee, K.B. Bang, S.B. Kim, I.H. Song, Development of tenofovir disoproxil fumarate resistance after complete viral suppression in a patient with treatment-naïve chronic hepatitis B: A case report and review of the literature, *World J. Gastroenterol.* 24 (2018) 1919–1924.
- [4] E.-S. Park, A.R. Lee, D.H. Kim, J.-H. Lee, J.-J. Yoo, S.H. Ahn, H. Sim, S. Park, H.S. Kang, J. Won, et al., Identification of a quadruple mutation that confers tenofovir resistance in chronic hepatitis B patients, *J. Hepatol.* 70 (2019) 1093–1102.
- [5] J. Mokaya, A.L. McNaughton, P.A. Bester, D. Goedhals, E. Barnes, B.D. Marsden, P.C. Matthews, Hepatitis B virus resistance to tenofovir: fact or fiction? A synthesis of the evidence to date, A systematic literature review and structural analysis of drug resistance mechanisms 5 (151) (2020), <https://doi.org/10.12688/wellcomeopenres.15992.1> Available at: <https://www.medrxiv.org/content/10.1101/19009563v1> [Accessed February 22, 2020].
- [6] M. Lemoine, S. Eholié, K. Lacombe, Reducing the neglected burden of viral hepatitis in Africa: strategies for a global approach, *J. Hepatol.* 62 (2015) 469–476.
- [7] J. Mokaya, A.L. McNaughton, M.J. Hadley, A. Beloukas, A.-M. Geretti, D. Goedhals, P.C. Matthews, A systematic review of hepatitis B virus (HBV) drug and vaccine escape mutations in Africa: a call for urgent action, *PLoS Negl. Trop. Dis.* 12 (2018) e0006629, <https://doi.org/10.1371/journal.pntd.0006629>.
- [8] World Health Organisation, Combating Hepatitis B and C to Reach Elimination by 2030, Available at: [http://apps.who.int/iris/bitstream/10665/206453/1/WHO\\_HIV\\_2016.04\\_eng.pdf?ua=1](http://apps.who.int/iris/bitstream/10665/206453/1/WHO_HIV_2016.04_eng.pdf?ua=1) [Accessed February 22, 2020] (2016).
- [9] G.S. Cooke, I. Andrieux-Meyer, T.L. Applegate, R. Atun, J.R. Burry, H. Cheinquer, G. Dusheiko, J.J. Feld, C. Gore, M.G. Griswold, et al., Accelerating the elimination of viral hepatitis: a Lancet Gastroenterology & Hepatology Commission, *Lancet Gastroenterol. Hepatol.* 4 (2019) 135–184.
- [10] World Health Organisation, Global Hepatitis Report, (2017) Available at: <https://www.who.int/hepatitis/publications/global-hepatitis-report2017/en/> [Accessed February 22, 2020].
- [11] T.G. Maponga, A.L. McNaughton, M. Van Schalkwyk, S. Hugo, C. Nwankwo, J. Taljaard, J. Mokaya, D.A. Smith, C. van Vuuren, D. Goedhals, et al., Treatment advantage in HBV/HIV coinfection compared to HBV monoinfection in a South African cohort, *J. Infect.* 81 (2020) 121–130.
- [12] C.W.N. Spearman, M.W. Sonderup, J.F. Botha, S.W. van der Merwe, E. Song, C. Kassianides, K.A. Newton, H.N. Hairwadzi, Division of Hepatology, Department of Medicine, University of Cape Town, South Africa, South African guideline for the management of chronic hepatitis B: 2013, *S. Afr. Med. J.* 103 (2013) 337–349.
- [13] World Health Organisation, Updated Recommendations on First-line and Second-line Antiretroviral Regimens and Post-exposure Prophylaxis and Recommendations on Early Infant Diagnosis of HIV, Available at: <http://www.who.int/hiv/pub/guidelines/ARV2018update/en/> [Accessed March 9, 2020] (2019).
- [14] A.L. McNaughton, H.E. Roberts, D. Bonsall, M. de Cesare, J. Mokaya, S.F. Lumley, T. Golubchik, P. Piazza, J.B. Martin, C. de Lara, et al., Illumina and Nanopore methods for whole genome sequencing of hepatitis B virus (HBV), *Sci. Rep.* 9 (2019) 7081.
- [15] N. Martel, S.A. Gomes, I. Chemin, C. Trépo, A. Kay, Improved rolling circle amplification (RCA) of hepatitis B virus (HBV) relaxed-circular serum DNA (RC-DNA), *J. Virol. Methods* 193 (2013) 653–659.
- [16] TANOT: a rapid BLAST-guided read mapper for small, divergent genomes. <https://bioinformatics.cvr.ac.uk/software/tanoti/>.
- [17] J.B. Singer, E.C. Thomson, J. McLauchlan, J. Hughes, R.J. Gifford, GLUE: a flexible software system for virus sequence data, *BMC Bioinformatics* 19 (2018) 532.
- [18] A. Beloukas, A.M. Geretti, Hepatitis B virus drug resistance, *Antimicrobial Drug Resistance*, Cham: Springer International Publishing, 2017.
- [19] A.L. McNaughton, P.A. Revill, M. Littlejohn, P.C. Matthews, M.A. Ansari, Analysis of genomic-length HBV sequences to determine genotype and subtype reference sequences, *J. Gen. Virol.* (2020), <https://doi.org/10.1099/jgv.0.001387>.
- [20] I.S. Min, C.H. Lee, I.S. Shin, N.E. Lee, H.S. Son, S.B. Kim, S.Y. Seo, S.B.H. Kim, S.B.W. Kim, S.O. Lee, et al., Treatment outcome and renal safety of 3-year tenofovir disoproxil fumarate therapy in chronic hepatitis B Patients with preserved glomerular filtration rate, *Gut Liver* 13 (2019) 93–103.
- [21] G. Cui, X. Xu, H. Diao, Comparative Meta-Analysis of Tenofovir Disoproxil Fumarate versus Emtricitabine and Tenofovir Disoproxil Fumarate as Treatments for Patients with Chronic Hepatitis B, *Sci. Rep.* 5 (2015) 1–11.
- [22] N.A. Louissaint, Y.-J. Cao, P.L. Skipper, R.G. Liberman, S.R. Tannenbaum, S. Nimmagadda, J.R. Anderson, S. Everts, R. Bakshi, E.J. Fuchs, et al., Single dose pharmacokinetics of oral tenofovir in plasma, peripheral blood mononuclear cells, colonic tissue, and vaginal tissue, *AIDS Res. Hum. Retroviruses* 29 (2013) 1443–1450.
- [23] C. Davis, G.S. Mgomella, A. da Silva Filipe, E.H. Frost, G. Giroux, J. Hughes, C. Hogan, P. Kaleebu, G. Asiki, J. McLauchlan, et al., Highly diverse hepatitis C strains detected in Sub-Saharan Africa have unknown susceptibility to direct-acting antiviral treatments, *Hepatology* 69 (2019) 1426–1441.
- [24] S. Faurati, C. Rodriguez, C. Hézode, A. Soulier, I. Ruiz, L. Poiteau, S. Chevaliez, J.-M. Pawlotsky, Frequent antiviral treatment failures in patients infected with hepatitis C virus genotype 4, subtype 4r, *Hepatology* 69 (2019) 513–523.